

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for TA347018

PKM2 (PKM) Mouse Monoclonal Antibody [Clone ID: 1F2-F4-H11]

Product data:

Product Type: Primary Antibodies

Clone Name: 1F2-F4-H11

Applications: IF, IP, WB

Recommend Dilution: WB: 1:1000

Reactivity: Human, Monkey

Host: Mouse Isotype: IgG2b

Clonality: Monoclonal

Immunogen: The immunogen for PKM2 antibody: purified recombinant human PKM2 protein fragments

expressed in E.coli.

Formulation: PBS(pH 7.4) containing with 0.03% Proclin300 and 50% glycerol.

Purification: Affinity purified

Predicted Protein Size: 60 kDa

Gene Name: pyruvate kinase, muscle

Database Link: NP 002645 Entrez Gene 697742 MonkeyEntrez Gene 5315 Human

Background: This gene encodes a protein involved in glycolysis. The encoded protein is a pyruvate kinase

that catalyzes the transfer of a phosphoryl group from phosphoenolpyruvate to ADP,

generating ATP and pyruvate. This protein has been shown to interact with thyroid hormone and may mediate cellular metabolic effects induced by thyroid hormones. This protein has been found to bind Opa protein, a bacterial outer membrane protein involved in gonococcal adherence to and invasion of human cells, suggesting a role of this protein in bacterial pathogenesis. Several alternatively spliced transcript variants encoding a few distinct

isoforms have been reported.

Synonyms: CTHBP; HEL-S-30; OIP3; PK3; PKM2; TCB; THBP1

Protein Families: Druggable Genome


Protein Pathways: Glycolysis / Gluconeogenesis, Metabolic pathways, Purine metabolism, Pyruvate metabolism,

Type II diabetes mellitus







Product images:



Western blot detection of PKM2 in HCT116, COS7, K562, A549 and Hela cell lysates using PKM2 mouse mAb (1:1000 diluted). Predicted band size:60KDa. Observed band size:60KDa.



Immunofluorescent analysis of 3T3 cells fixed by anhydrous methanol for 2h at -20? and using anti-PKM2 mouse mAb (dilution 1:500).



Immunoprecipitation analysis of Hela cell lysates using PKM2.