

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# **Product datasheet for TA500939**

## FAHD2A Mouse Monoclonal Antibody [Clone ID: OTI6D9]

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI6D9

**Applications:** FC, IF, IHC, WB

**Recommend Dilution:** WB 1:2000, IHC 1:50, IF 1:100, Flow 1:100

Reactivity: Human
Host: Mouse
Isotype: IgG2b

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human FAHD2A (NP\_057128) produced in

HEK293T cell.

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

**Concentration:** 0.69 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

**Predicted Protein Size:** 34.6 kDa

**Gene Name:** fumarylacetoacetate hydrolase domain containing 2A


Database Link: NP 057128 Entrez Gene 51011 Human

**Background:** May have hydrolase activity


Synonyms: CGI-105



### **Product images:**




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY FAHD2A ([RC211128], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-FAHD2A. Positive lysates [LY402492] (100ug) and [LC402492] (20ug) can be purchased separately from OriGene.




Immunohistochemical staining of paraffinembedded Kidney tissue within the normal limits using anti-FAHD2Amouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA500939, Dilution 1:50)







Immunohistochemical staining of paraffinembedded liver tissue within the normal limits using anti-FAHD2Amouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA500939, Dilution 1:50)




Immunohistochemical staining of paraffinembedded Carcinoma of liver tissue using anti-FAHD2Amouse monoclonal antibody. (Heatinduced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA500939, Dilution 1:50)



Anti-FAHD2A mouse monoclonal antibody (TA500939) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY FAHD2A ([RC211128]).





HEK293T cells transfected with either pCMV6-ENTRY FAHD2A ([RC211128]) (Red) or empty vector control plasmid (Blue) were immunostained with anti-FAHD2A mouse monoclonal (TA500939), and then analyzed by flow cytometry.

Flow cytometric analysis of Jurkat cells, using anti-FAHD2A antibody (TA500939), (Red) compared to a nonspecific negative control antibody ([TA50011]) (Blue).