

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## **Product datasheet for TA501648**

## Protein Kinase D2 (PRKD2) Mouse Monoclonal Antibody [Clone ID: OTI5A3]

**Product data:** 

**Product Type:** Primary Antibodies

Clone Name: OTI5A3

**Applications:** FC, IF, IHC, WB

**Recommend Dilution:** WB 1:2000, IHC 1:150, IF 1:100, FLOW 1:100

Reactivity: Human
Host: Mouse
Isotype: IgG2b

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human PRKD2 (NP\_057541) produced in HEK293T

cell.

**Formulation:** PBS (PH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

**Concentration:** 0.97 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

**Predicted Protein Size:** 96.5 kDa

**Gene Name:** protein kinase D2

Database Link: NP 057541 Entrez Gene 25865 Human

**Background:** The protein encoded by this gene belongs to the protein kinase D (PKD) family of

serine/threonine protein kinases. This kinase can be activated by phorbol esters as well as by

gastrin via the cholecystokinin B receptor (CCKBR) in gastric cancer cells. It can bind to diacylglycerol (DAG) in the trans-Golgi network (TGN) and may regulate basolateral

membrane protein exit from TGN. Alternative splicing results in multiple transcript variants

encoding different isoforms. [provided by RefSeq]


**Synonyms:** HSPC187; nPKC-D2; PKD2

**Protein Families:** Druggable Genome, Protein Kinase







## **Product images:**



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY PRKD2 ([RC215335], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-PRKD2. Positive lysates [LY402555] (100ug) and [LC402555] (20ug) can be purchased separately from OriGene.




Immunohistochemical staining of paraffinembedded Carcinoma of Human bladder tissue using anti-PRKD2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA501648)




Anti-PRKD2 mouse monoclonal antibody (TA501648) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY PRKD2 ([RC215335]).






HEK293T cells transfected with either [RC215335] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-PRKD2 antibody (TA501648), and then analyzed by flow cytometry.



Flow cytometric Analysis of Hela cells, using anti-PRKD2 antibody (TA501648), (Red), compared to a nonspecific negative control antibody ([TA50011]), (Blue).



Flow cytometric Analysis of Jurkat cells, using anti-PRKD2 antibody (TA501648), (Red), compared to a nonspecific negative control antibody ([TA50011]), (Blue).