

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# **Product datasheet for TA501749**

## **ARHGAP25 Mouse Monoclonal Antibody [Clone ID: OTI3G1]**

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI3G1

**Applications:** FC, IF, IHC, WB

**Recommend Dilution:** WB 1:500~2000, IHC 1:150, IF 1:100, FLOW 1:100

Reactivity: Human
Host: Mouse
Isotype: IgG2a

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human ARHGAP25 (NP\_055697) produced in

HEK293T cell.

**Formulation:** PBS (PH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

**Concentration:** 0.78 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Predicted Protein Size: 72.3 kDa

Gene Name: Rho GTPase activating protein 25

Database Link: NP 055697 Entrez Gene 9938 Human

Background: ARHGAPs, such as ARHGAP25, encode negative regulators of Rho GTPases (see ARHA; MIM

165390), which are implicated in actin remodeling, cell polarity, and cell migration (Katoh and


Katoh, 2004 [PubMed 15254788]). [supplied by OMIM]

Synonyms: HEL-S-308; KAIA0053






### **Product images:**




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY ARHGAP25 ([RC217414], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-ARHGAP25. Positive lysates [LY402385] (100ug) and [LC402385] (20ug) can be purchased separately from OriGene.




Western blot analysis of extracts (35ug) from 9 different cell lines by using anti-ARHGAP25 monoclonal antibody.






Immunohistochemical staining of paraffinembedded Human lymph node tissue within the normal limits using anti-ARHGAP25 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA501749, Dilution 1:50)




Immunohistochemical staining of paraffinembedded Human lymphoma tissue using anti-ARHGAP25 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA501749, Dilution 1:50)



Anti-ARHGAP25 mouse monoclonal antibody (TA501749) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY ARHGAP25 ([RC217414]).





HEK293T cells transfected with either [RC217414] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-ARHGAP25 antibody (TA501749), and then analyzed by flow cytometry.