

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for TA501760

PIK3AP1 Mouse Monoclonal Antibody [Clone ID: OTI7B12]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI7B12

Applications: FC, IF, IHC, WB

Recommend Dilution: WB 1:2000, IHC 1:150, IF 1:100, FLOW 1:100

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human PIK3AP1 (NP_689552) produced in

HEK293T cell.

Formulation: PBS (PH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 0.4 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Predicted Protein Size: 90.2 kDa

Gene Name: phosphoinositide-3-kinase adaptor protein 1

Database Link: NP 689522 Entrez Gene 118788 Human

Background: Involved in the activation of phosphoinositide 3-kinase (PI3K) in B-cells and in natural killer

(NK) cells. Couples B-cell antigen receptor (BCR) to PI3K activation by providing a docking site for the PI3K subunit PIK3R1, which contributes to B-cell development. Seems to have a complementary role with CD19 in PI3K activation (By similarity). May be involved in the

survival of mature B cells via activation of REL (By similarity).

Synonyms: BCAP


Protein Families: Druggable Genome

Protein Pathways: B cell receptor signaling pathway







Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY PIK3AP1 ([RC214125], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-PIK3AP1. Positive lysates [LY403461] (100ug) and [LC403461] (20ug) can be purchased separately from OriGene.




Immunohistochemical staining of paraffinembedded Human lymph node tissue within the normal limits using anti-PIK3AP1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA501760, Dilution 1:50)



Anti-PIK3AP1 mouse monoclonal antibody (TA501760) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY PIK3AP1 ([RC214125]).





HEK293T cells transfected with either [RC214125] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-PIK3AP1 antibody (TA501760), and then analyzed by flow cytometry.