

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# **Product datasheet for TA504303**

## PSMD3 Mouse Monoclonal Antibody [Clone ID: OTI2C2]

### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI2C2
Applications: FC, IF, WB

Recommend Dilution: WB 1:2000, IF 1:100, FLOW 1:100

Reactivity: Human
Host: Mouse
Isotype: IgG2a

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human PSMD3(NP\_002800) produced in HEK293T

cell

**Formulation:** PBS (PH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Predicted Protein Size: 60.8 kDa

Gene Name: proteasome 26S subunit, non-ATPase 3

Database Link: NP 002800 Entrez Gene 5709 Human

**Background:** The 26S proteasome is a multicatalytic proteinase complex with a highly ordered structure

composed of 2 complexes, a 20S core and a 19S regulator. The 20S core is composed of 4 rings of 28 non-identical subunits; 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. The 19S regulator is composed of a base, which contains 6 ATPase subunits and 2 non-ATPase subunits, and a lid, which contains up to 10 non-ATPase subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. An essential function of a modified proteasome, the immunoproteasome, is the processing of class I MHC peptides. This gene encodes one of the non-ATPase subunits of the 19S regulator

lid. [provided by RefSeq, Jul 2008]


**Synonyms:** P58; RPN3; S3; TSTA2






**Protein Pathways:** Proteasome


## **Product images:**



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY PSMD3 ([RC202307], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-PSMD3. Positive lysates [LY400996] (100ug) and [LC400996] (20ug) can be purchased separately from OriGene.




Anti-PSMD3 mouse monoclonal antibody (TA504303) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY PSMD3 ([RC202307]).



HEK293T cells transfected with either [RC202307] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-PSMD3 antibody (TA504303), and then analyzed by flow cytometry.





Flow cytometric Analysis of Jurkat cells, using anti-PSMD3 antibody (TA504303), (Red), compared to a nonspecific negative control antibody, (Blue).