

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for TA505025

C1ORF151 (MINOS1) Mouse Monoclonal Antibody [Clone ID: OTI4H7]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI4H7
Applications: IHC, WB

Recommend Dilution: WB 1:1000, IHC 1:150

Reactivity: Human
Host: Mouse
Isotype: IgG2b

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human C1orf151(NP_001007535) produced in

HEK293T cell.

Formulation: PBS (PH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Predicted Protein Size: 8.6 kDa

Gene Name: mitochondrial contact site and cristae organizing system subunit 10


Database Link: NP 001027535 Entrez Gene 440574 Human

Synonyms: C1orf151; Mic10; MIO10


Protein Families: Transmembrane



Product images:




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY C1orf151 ([RC212930], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-C1orf151. Positive lysates [LY422307] (100ug) and [LC422307] (20ug) can be purchased separately from OriGene.




Western blot analysis of extracts (10ug) from 5 diff1 erent cell lines by using anti-C1orf151 monoclonal antibody (1:200).






Immunohistochemical staining of paraffinembedded Human breast tissue within the normal limits using anti-C1orf151 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, TA505025)




Immunohistochemical staining of paraffinembedded Adenocarcinoma of Human breast tissue using anti-C1orf151 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, TA505025)




Immunohistochemical staining of paraffinembedded Human colon tissue within the normal limits using anti-C1orf151 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, TA505025)






Immunohistochemical staining of paraffinembedded Human Kidney tissue within the normal limits using anti-C1orf151 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, TA505025)





Immunohistochemical staining of paraffinembedded Human liver tissue within the normal limits using anti-C1orf151 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, TA505025)




Immunohistochemical staining of paraffinembedded Carcinoma of Human liver tissue using anti-C1orf151 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, TA505025)









Immunohistochemical staining of paraffinembedded Adenocarcinoma of Human endometrium tissue using anti-C1orf151 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, TA505025)