

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# **Product datasheet for TA506337**

# **CD7 Mouse Monoclonal Antibody [Clone ID: OTI1A6]**

### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI1A6
Applications: IHC, WB

Recommend Dilution: WB 1:2000, IHC 1:150

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human CD7(NP\_006128) produced in HEK293T

cell.

**Formulation:** PBS (PH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Predicted Protein Size: 22.9 kDa

**Gene Name:** CD7 molecule

Database Link: NP 006128 Entrez Gene 924 Human

**Background:** This gene encodes a transmembrane protein which is a member of the immunoglobulin

superfamily. This protein is found on thymocytes and mature T cells. It plays an essential role in T-cell interactions and also in T-cell/B-cell interaction during early lymphoid development.

[provided by RefSeq, Jul 2008]

Synonyms: GP40; LEU-9; Tp40; TP41


**Protein Families:** Druggable Genome, Transmembrane

Protein Pathways: Hematopoietic cell lineage







## **Product images:**



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY CD7 ([RC201231], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-CD7. Positive lysates [LY416844] (100ug) and [LC416844] (20ug) can be purchased separately from OriGene.



Western blot analysis of extracts (10ug) from 1 cell line by using anti-CD7 monoclonal antibody (1:200).



Immunohistochemical staining of paraffinembedded Human lymph node tissue within the normal limits using anti-CD7 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, TA506337)