

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for TA506533

alpha Synuclein (SNCA) Mouse Monoclonal Antibody [Clone ID: OTI2F6]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI2F6
Applications: IHC, WB

Recommend Dilution: WB 1:4000, IHC: 1:150

Reactivity: Human
Host: Mouse
Isotype: IgG2b

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human SNCA(NP_000336) produced in HEK293T

cell.

Formulation: PBS (PH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Predicted Protein Size: 14.3 kDa

Gene Name: synuclein alpha

Database Link: NP 000336 Entrez Gene 6622 Human

Synonyms: NACP; PARK1; PARK4; PD1


Protein Families: Druggable Genome

Protein Pathways: Alzheimer's disease, Parkinson's disease






Product images:




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY SNCA ([RC210606], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-SNCA. Positive lysates [LY400127] (100ug) and [LC400127] (20ug) can be purchased separately from OriGene.




Immunohistochemical staining of paraffinembedded Adenocarcinoma of Human colon tissue using anti-SNCA mouse monoclonal antibody. (TA506533) (1:150).





Immunohistochemical staining of paraffinembedded Human adult brain tissue within the normal limits using anti-SNCA mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, TA506533) (1:500)



Immunohistochemical staining of paraffinembedded Human embryonic cerebellum within the normal limits using anti-SNCA mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, TA506533) (1:500)