

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## **Product datasheet for TA506752**

## **ARG2 Mouse Monoclonal Antibody [Clone ID: OTI1F6]**

**Product data:** 

**Product Type:** Primary Antibodies

Clone Name: OTI1F6

**Applications:** WB

Recommend Dilution: WB 1:4000

Reactivity: Human Host: Mouse

**Isotype:** IgG2b

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human ARG2(NP\_001163) produced in HEK293T

cell

**Formulation:** PBS (PH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

**Gene Name:** arginase 2

Database Link: NP 001163 Entrez Gene 384 Human

**Background:** Arginase catalyzes the hydrolysis of arginine to ornithine and urea. At least two isoforms of

mammalian arginase exists (types I and II) which differ in their tissue distribution, subcellular

localization, immunologic crossreactivity and physiologic function. The type II isoform

encoded by this gene, is located in the mitochondria and expressed in extra-hepatic tissues, especially kidney. The physiologic role of this isoform is poorly understood; it is thought to play a role in nitric oxide and polyamine metabolism. Transcript variants of the type II gene resulting from the use of alternative polyadenylation sites have been described. [provided by

RefSeq, Jul 2008]

**Synonyms:** arginase; arginase 2; kidney arginase; L-arginine amidinohydrolase; L-arginine ureahydrolase;


nonhepatic arginase; type II

**Protein Pathways:** Arginine and proline metabolism, Metabolic pathways





## **Product images:**



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY ARG2 ([RC206756], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-ARG2. Positive lysates [LY420091] (100ug) and [LC420091] (20ug) can be purchased separately from OriGene.