

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for TA800385

CD20 (MS4A1) Mouse Monoclonal Antibody [Clone ID: OTI4B4]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI4B4
Applications: IHC, WB

Recommend Dilution: WB 1:2000, IHC 1:150

Reactivity: Human, Monkey, Mouse, Dog

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human MS4A1 (NP_068769) produced in HEK293T

cell

Formulation: PBS (PH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Predicted Protein Size: 32.9 kDa

Gene Name: membrane spanning 4-domains A1

Database Link: NP 068769 Entrez Gene 12482 MouseEntrez Gene 485430 DogEntrez Gene 696843

MonkeyEntrez Gene 931 Human

Background: This gene encodes a member of the membrane-spanning 4A gene family. Members of this

nascent protein family are characterized by common structural features and similar

intron/exon splice boundaries and display unique expression patterns among hematopoietic cells and nonlymphoid tissues. This gene encodes a B-lymphocyte surface molecule which plays a role in the development and differentiation of B-cells into plasma cells. This family member is localized to 11q12, among a cluster of family members. Alternative splicing of this gene results in two transcript variants which encode the same protein. [provided by RefSeq,

Jul

Synonyms: B1; Bp35; CD20; CVID5; LEU-16; MS4A2; S7


Protein Families: Druggable Genome, Transmembrane







Protein Pathways: Hematopoietic cell lineage

Product images:







HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY MS4A1 ([RC201242], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-MS4A1. Positive lysates [LY402889] (100ug) and [LC402889] (20ug) can be purchased separately from OriGene.

Western blot analysis of extracts (35ug) from 9 different cell lines by using anti-MS4A1 monoclonal antibody (HepG2: human; HeLa: human; SVT2: mouse; A549: human; COS7: monkey; Jurkat: human; MDCK: canine; PC12: rat; MCF7: human).

Immunohistochemical staining of paraffinembedded Human tonsil within the normal limits using anti-MS4A1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 100°C for 10min, TA800385)