

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for TA800714

Cytokeratin 5 (KRT5) Mouse Monoclonal Antibody [Clone ID: OTI1F8]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI1F8

Applications: IHC

Reactivity: Human

Host: Mouse

Isotype: IgG1

Clonality: Monoclonal

Immunogen: Synthetic peptide: EQGTKTVRQN LEPLFEQYIN NLRRQ, corresponding to the conserved region

of human KRT5 and KRT6

Formulation: PBS (PH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Predicted Protein Size: 62.2 kDa

Gene Name: keratin 5

Database Link: NP 000415 Entrez Gene 3852 Human

Background: The protein encoded by this gene is a member of the keratin gene family. The type II

cytokeratins consist of basic or neutral proteins which are arranged in pairs of heterotypic keratin chains coexpressed during differentiation of simple and stratified epithelial tissues. This type II cytokeratin is specifically expressed in the basal layer of the epidermis with family member KRT14. Mutations in these genes have been associated with a complex of diseases termed epidermolysis bullosa simplex. The type II cytokeratins are clustered in a region of


chromosome 12q12-q13. [provided by RefSeq, Jul 2008]

Synonyms: CK5; DDD; DDD1; EBS2; K5; KRT5A







Product images:



Immunohistochemical staining of paraffinembedded Human breast tissue within the normal limits using anti-KRT5 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, TA800714)




Immunohistochemical staining of paraffinembedded Adenocarcinoma of Human colon tissue using anti-KRT5 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, TA800714)



Immunohistochemical staining of paraffinembedded Human prostate tissue within the normal limits using anti-KRT5 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, TA800714)





Immunohistochemical staining of paraffinembedded Carcinoma of Human prostate tissue using anti-KRT5 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, TA800714)