

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# Product datasheet for TA804188

# GNG7 Mouse Monoclonal Antibody [Clone ID: OTI3G1]

## **Product data:**

Product Type:	Primary Antibodies		
Clone Name:	OTI3G1		
Applications:	WB		
<b>Recommend Dilution:</b>	WB 1:2000		
Reactivity:	Human		
Host:	Mouse		
lsotype:	lgG2a		
Clonality:	Monoclonal		
Immunogen:	Human recombinant protein fragment corresponding to amino acids 2-65 of human GNG7(NP_443079) produced in E.coli.		
Formulation:	PBS (PH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.		
Concentration:	1 mg/ml		
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)		
Predicted Protein Size:	7.3 kDa		
Gene Name:	G protein subunit gamma 7		
Database Link:	<u>NP_443079 Entrez Gene 2788 Human</u>		
Background:	Guanine nucleotide-binding proteins (G proteins) are involved as a modulator or transducer in various transmembrane signaling systems. The beta and gamma chains are required for the GTPase activity, for replacement of GDP by GTP, and for G protein-effector interaction. Plays a role in the regulation of adenylyl cyclase signaling in certain regions of the brain. Plays a role in the formation or stabilzation of a G protein heterotrimer (G(olf) subunit alpha-beta- gamma-7) that is required for adenylyl cyclase activity in the striatum. [UniProtKB/Swiss-Prot Function]		
Synonyms:	FLJ00058		
Protein Families:	Druggable Genome		
Protein Pathways:	Chemokine signaling pathway		



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2020 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

### **Product images:**

170	_	
130	_	
100	_	
70	—	
55		
40		
35	—	
25		
15	—	
10	—	-

HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY GNG7 ([RC208768], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-GNG7 (1:2000). Positive lysates [LY409457] (100ug) and [LC409457] (20ug) can be purchased separately from OriGene.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2020 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US