

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## **Product datasheet for TA804327**

## XLF (NHEJ1) Mouse Monoclonal Antibody [Clone ID: OTI1B5]

**Product data:** 

**Product Type:** Primary Antibodies

Clone Name: OTI1B5
Applications: IHC, WB

Recommend Dilution: WB 1:2000, IHC 1:150

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human NHEJ1 (NP\_079058) produced in HEK293T

cell

**Formulation:** PBS (PH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Predicted Protein Size: 33.2 kDa

Gene Name: non-homologous end joining factor 1

Database Link: NP 079058 Entrez Gene 79840 Human

**Background:** Double-strand breaks in DNA result from genotoxic stresses and are among the most

damaging of DNA lesions. This gene encodes a DNA repair factor essential for the nonhomologous end-joining pathway, which preferentially mediates repair of double-stranded breaks. Mutations in this gene cause different kinds of severe combined

immunodeficiency disorders. [provided by RefSeq, Jul


Synonyms: XLF

**Protein Pathways:** Non-homologous end-joining






## **Product images:**




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY NHEJ1 ([RC203393], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-NHEJ1. Positive lysates [LY403031] (100ug) and [LC403031] (20ug) can be purchased separately from OriGene.




Equivalent amounts of cell lysates (10 ug per lane) of wild-type HeLa cells (WT, Cat# LC810HELA) and NHEJ1-Knockout HeLa cells (KO, Cat# [LC831412]) were separated by SDS-PAGE and immunoblotted with anti-NHEJ1 monoclonal antibody TA804327 (1:100). Then the blotted membrane was stripped and reprobed with anti-HSP90 antibody as a loading control.






Immunohistochemical staining of paraffinembedded Carcinoma of Human liver tissue using anti-NHEJ1 Mouse monoclonal antibody. (TA804327; heat-induced epitope retrieval by 1 mM EDTA in 10mM Tris, pH8.5, 120°C for 3min)



Immunohistochemical staining of paraffinembedded Carcinoma of Human lung tissue using anti-NHEJ1 Mouse monoclonal antibody. (TA804327; heat-induced epitope retrieval by 1 mM EDTA in 10mM Tris, pH8.5, 120°C for 3min)



Immunohistochemical staining of paraffinembedded Carcinoma of Human pancreas tissue using anti-NHEJ1 Mouse monoclonal antibody. (TA804327; heat-induced epitope retrieval by 1 mM EDTA in 10mM Tris, pH8.5, 120°C for 3min)