

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# **Product datasheet for TA804390**

### **Eotaxin (CCL11) Mouse Monoclonal Antibody [Clone ID: OTI2H9]**

#### **Product data:**

**Product Type: Primary Antibodies** 

Clone Name: OTI2H9

**Applications:** WB

**Recommend Dilution:** WB 1:2000

Reactivity: Human

Host: Isotype: lgG1

Clonality: Monoclonal

Immunogen: Human recombinant protein fragment corresponding to amino acids 24-97 of human CCL11

(NP 002977) produced in E.coli.

Formulation: PBS (PH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Mouse

**Predicted Protein Size:** 8.3 kDa

Gene Name: C-C motif chemokine ligand 11

NP 002977 Entrez Gene 6356 Human Database Link:

Background: This gene is one of several chemokine genes clustered on the q-arm of chromosome 17.

> Chemokines form a superfamily of secreted proteins involved in immunoregulatory and inflammatory processes. The superfamily is divided into four subfamilies based on the arrangement of the N-terminal cysteine residues of the mature peptide. This chemokine, a

member of the CC subfamily, displays chemotactic activity for eosinophils, but not mononuclear cells or neutrophils. This eosinophil-specific chemokine is thought to be involved in eosinophilic inflammatory diseases such as atopic dermatitis, allergic rhinitis,

asthma and parasitic infections. [provided by RefSeq, Jul 2013]

SCYA11 Synonyms:

**Protein Families:** Druggable Genome, Secreted Protein, Transmembrane






**Protein Pathways:** 

Asthma, Chemokine signaling pathway, Cytokine-cytokine receptor interaction, NOD-like receptor signaling pathway

## **Product images:**



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY CCL11 ([RC204726], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-CCL11. Positive lysates [LY401045] (100ug) and [LC401045] (20ug) can be purchased separately from OriGene.