

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for TA804739

MOX1 (MEOX1) Mouse Monoclonal Antibody [Clone ID: OTI5B11]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI5B11
Applications: IHC, WB

Recommend Dilution: WB 1:2000, IHC 1:150

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human MEOX1 (NP_004518) produced in E.coli.

Formulation: PBS (PH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Predicted Protein Size: 27.8 kDa

Gene Name: mesenchyme homeobox 1

Database Link: NP 004518 Entrez Gene 4222 Human

Background: This gene encodes a member of a subfamily of non-clustered, diverged, antennapedia-like

homeobox-containing genes. The encoded protein may play a role in the molecular signaling network regulating somite development. Alternatively spliced transcript variants encoding

Hetwork regulating somite development. Atternatively spliced transcript variants end

different isoforms have been described. [provided by RefSeq, Jul 2008]


Synonyms: KFS2; MOX1

Protein Families: Transcription Factors






Product images:




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY MEOX1 ([RC210106], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-MEOX1. Positive lysates [LY417902] (100ug) and [LC417902] (20ug) can be purchased separately from OriGene.



Immunohistochemical staining of paraffinembedded Human lymph node tissue within the normal limits using anti-MEOX1 Mouse monoclonal antibody. (TA804739; heat-induced epitope retrieval by 1 mM EDTA in 10mM Tris, pH8.5, 120°C for 3min)





Immunohistochemical staining of paraffinembedded Human tonsil within the normal limits using anti-MEOX1 Mouse monoclonal antibody. (TA804739; heat-induced epitope retrieval by 1 mM EDTA in 10mM Tris, pH8.5, 120°C for 3min)