

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# **Product datasheet for TA804783**

## **CLOCK Mouse Monoclonal Antibody [Clone ID: OTI2H7]**

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI2H7
Applications: IHC, WB

Recommend Dilution: WB 1:2000, IHC 1:150

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

Immunogen: Human recombinant protein fragment corresponding to amino acids 1-285 of human CLOCK

(NP\_004889) produced in E.coli.

**Formulation:** PBS (PH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

**Predicted Protein Size:** 95.1 kDa

**Gene Name:** clock circadian regulator

Database Link: NP 004889 Entrez Gene 9575 Human


Synonyms: bHLHe8; KAT13D

**Protein Families:** Druggable Genome, Transcription Factors


**Protein Pathways:** Circadian rhythm - mammal



# **Product images:**




Human recombinant protein fragment corresponding to amino acids 1-285 of human CLOCK (NP\_004889) produced in E.coli.




Immunohistochemical staining of paraffinembedded Carcinoma of Human pancreas tissue using anti-CLOCK mouse monoclonal antibody. (Heat-induced epitope retrieval by 1 mM EDTA in 10mM Tris, pH8.5, 120°C for 3min, TA804783)






Immunohistochemical staining of paraffinembedded Human bladder tissue within the normal limits using anti-CLOCK mouse monoclonal antibody. (Heat-induced epitope retrieval by 1 mM EDTA in 10mM Tris, pH8.5, 120°C for 3min, TA804783)



Immunohistochemical staining of paraffinembedded Carcinoma of Human bladder tissue using anti-CLOCK mouse monoclonal antibody. (Heat-induced epitope retrieval by 1 mM EDTA in 10mM Tris, pH8.5, 120°C for 3min, TA804783)



Immunohistochemical staining of paraffinembedded Human tonsil within the normal limits using anti-CLOCK mouse monoclonal antibody. (Heat-induced epitope retrieval by 1 mM EDTA in 10mM Tris, pH8.5, 120°C for 3min, TA804783)