

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for TA804816

R Cadherin (CDH4) Mouse Monoclonal Antibody [Clone ID: OTI6H5]

Product data:

Product Type: Primary Antibodies

Clone Name: **OTI6H5** IHC. WB **Applications:**

Recommend Dilution: WB 1:2000, IHC 1:150

Reactivity: Human Mouse Host: Isotype: lgG1

Clonality: Monoclonal

Immunogen: Human recombinant protein fragment corresponding to amino acids 170-428 of human

CDH4 (NP 001785) produced in E.coli.

Formulation: PBS (PH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Predicted Protein Size: 98.3 kDa Gene Name: cadherin 4

Database Link: NP 001785 Entrez Gene 1002 Human

Background: This gene is a classical cadherin from the cadherin superfamily. The encoded protein is a

calcium-dependent cell-cell adhesion glycoprotein comprised of five extracellular cadherin repeats, a transmembrane region and a highly conserved cytoplasmic tail. Based on studies

in chicken and mouse, this cadherin is thought to play an important role during brain

segmentation and neuronal outgrowth. In addition, a role in kidney and muscle development is indicated. Of particular interest are studies showing stable cis-heterodimers of cadherins 2 and 4 in cotransfected cell lines. Previously thought to interact in an exclusively homophilic manner, this is the first evidence of cadherin heterodimerization. Three transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Nov 2011]


Synonyms: CAD4; R-CAD; RCAD **Protein Families:** Transmembrane






Protein Pathways: Cell adhesion molecules (CAMs)

Product images:




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY CDH4 ([RC210970], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-CDH4. Positive lysates [LY400684] (100ug) and [LC400684] (20ug) can be purchased separately from OriGene.




Immunohistochemical staining of paraffinembedded Carcinoma of Human thyroid tissue using anti-CDH4 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1 mM EDTA in 10mM Tris, pH8.5, 120°C for 3min, TA804816)





Immunohistochemical staining of paraffinembedded Carcinoma of Human kidney tissue using anti-CDH4 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1 mM EDTA in 10mM Tris, pH8.5, 120°C for 3min, TA804816)



Immunohistochemical staining of paraffinembedded Human Kidney tissue within the normal limits using anti-CDH4 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1 mM EDTA in 10mM Tris, pH8.5, 120°C for 3min, TA804816)