

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for TA806789

Dishevelled 2 (DVL2) Mouse Monoclonal Antibody [Clone ID: OTI7C7]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI7C7
Applications: IHC, WB

Recommend Dilution: WB 1:2000, IHC 1:250

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

Immunogen: Human recombinant protein fragment corresponding to amino acids 1-320 of human

DVL2(NP_004413) produced in E.coli.

Formulation: PBS (PH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Predicted Protein Size: 78.8 kDa

Gene Name: dishevelled segment polarity protein 2

Database Link: NP 004413 Entrez Gene 1856 Human

Background: This gene encodes a member of the dishevelled (dsh) protein family. The vertebrate dsh

proteins have approximately 40% amino acid sequence similarity with Drosophila dsh. This gene encodes a 90-kD protein that undergoes posttranslational phosphorylation to form a 95-kD cytoplasmic protein, which may play a role in the signal transduction pathway

mediated by multiple Wnt proteins. The mechanisms of dishevelled function in Wnt signaling

are likely to be conserved among metazoans. [provided by RefSeq, Jul 2008]

Synonyms: dishevelled; dishevelled 2; dishevelled 2 (homologous to Drosophila dsh); DSH homolog 2;

dsh homolog 2 (Drosophila); segment polarity protein dishevelled homolog DVL-2

Protein Families: Druggable Genome, ES Cell Differentiation/IPS


Protein Pathways: Basal cell carcinoma, Colorectal cancer, Melanogenesis, Notch signaling pathway, Pathways in

cancer, Wnt signaling pathway






Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY DVL2 ([RC202233], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-DVL2. Positive lysates [LY401405] (100ug) and [LC401405] (20ug) can be purchased separately from OriGene.



Immunohistochemical staining of paraffinembedded Adenocarcinoma of Human endometrium tissue using anti-DVL2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1 mM EDTA in 10mM Tris, pH8.5, 120°C for 3min, TA806789)