

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## **Product datasheet for TA807497**

## SFRP2 Mouse Monoclonal Antibody [Clone ID: OTI6C2]

**Product data:** 

**Product Type:** Primary Antibodies

Clone Name: OTI6C2

Applications: IHC, WB

Recommend Dilution: IHC 1:150

Reactivity: Human

Host: Mouse

Isotype: IgG2a

Clonality: Monoclonal

**Immunogen:** Human recombinant protein fragment corresponding to amino acids 25-295 of human

SFRP2(NP\_003004) produced in E.coli.

**Formulation:** PBS (PH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Predicted Protein Size: 31.4 kDa

Gene Name: secreted frizzled related protein 2

Database Link: NP 003004 Entrez Gene 6423 Human

**Background:** This gene encodes a member of the SFRP family that contains a cysteine-rich domain

homologous to the putative Wnt-binding site of Frizzled proteins. SFRPs act as soluble modulators of Wnt signaling. Methylation of this gene is a potential marker for the presence

of colorectal cancer. [provided by RefSeq, Jul 2008]


Synonyms: FRP-2; SARP1; SDF-5
Protein Families: Secreted Protein

**Protein Pathways:** Wnt signaling pathway






## **Product images:**




Western blot analysis of extracts (35ug) from 3 different cell lines and 5 different tissue lysates by using anti-SFRP2 monoclonal antibody (1:500).




Immunohistochemical staining of paraffinembedded Human Kidney tissue within the normal limits using anti-SFRP2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1 mM EDTA in 10mM Tris, pH8.5, 120°C for 3min, TA807497)






Immunohistochemical staining of paraffinembedded Human liver tissue within the normal limits using anti-SFRP2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1 mM EDTA in 10mM Tris, pH8.5, 120°C for 3min, TA807497)




Immunohistochemical staining of paraffinembedded Carcinoma of Human liver tissue using anti-SFRP2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1 mM EDTA in 10mM Tris, pH8.5, 120°C for 3min, TA807497)



Immunohistochemical staining of paraffinembedded Carcinoma of Human lung tissue using anti-SFRP2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1 mM EDTA in 10mM Tris, pH8.5, 120°C for 3min, TA807497)





Immunohistochemical staining of paraffinembedded Adenocarcinoma of Human ovary tissue using anti-SFRP2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1 mM EDTA in 10mM Tris, pH8.5, 120°C for 3min, TA807497)