

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# **Product datasheet for TA807631**

## **IKBKE Mouse Monoclonal Antibody [Clone ID: OTI1E4]**

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI1E4
Applications: IHC, WB

**Recommend Dilution:** WB 1:500, IHC 1:150

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

**Immunogen:** Human recombinant protein fragment corresponding to amino acids 474-716 of human

IKBKE(NP\_054721) produced in E.coli.

**Formulation:** PBS (PH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Predicted Protein Size: 80.3 kDa

**Gene Name:** inhibitor of kappa light polypeptide gene enhancer in B-cells, kinase epsilon

Database Link: NP 054721 Entrez Gene 9641 Human

Background: IKBKE is a noncanonical I-kappa-B (see MIM 164008) kinase (IKK) that is essential for

regulating antiviral signaling pathways. IKBKE has also been identified as a breast cancer

(MIM 114480) oncogene and is amplified and overexpressed in over 30% of breast

carcinomas and breast cancer cell lines (Hutti et al., 2009 [PubMed 19481526]). [supplied by

OMIM, Oct 2009]

Synonyms: IKK-E; IKK-i; IKKE; IKKI

**Protein Families:** Druggable Genome, Protein Kinase


**Protein Pathways:** Cytosolic DNA-sensing pathway, RIG-I-like receptor signaling pathway, Toll-like receptor

signaling pathway






## **Product images:**



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY IKBKE ([RC212481], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-IKBKE (1:500). Positive lysates [LY415533] (100ug) and [LC415533] (20ug) can be purchased separately from OriGene.



Immunohistochemical staining of paraffinembedded Human lung tissue within the normal limits using anti-IKBKE mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, TA807631) (1:150)