

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for TA807749

PODXL Mouse Monoclonal Antibody [Clone ID: OTI8D6]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI8D6

Applications: IHC

Recommend Dilution: IHC 1:150

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

Immunogen: Human recombinant protein fragment corresponding to amino acids 32-461 of human

PODXL(NP_005388) produced in HEK293 cells.

Formulation: PBS (PH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Gene Name: podocalyxin like

Database Link: NP 001018121 Entrez Gene 5420 Human

Background: This gene encodes a member of the sialomucin protein family. The encoded protein was

originally identified as an important component of glomerular podocytes. Podocytes are highly differentiated epithelial cells with interdigitating foot processes covering the outer aspect of the glomerular basement membrane. Other biological activities of the encoded protein include: binding in a membrane protein complex with Na+/H+ exchanger regulatory

factor to intracellular cytoskeletal elements, playing a role in hematopoetic cell

differentiation, and being expressed in vascular endothelium cells and binding to L-selectin.

[provided by RefSeq, Jul 2008]


Synonyms: Gp200; PC; PCLP; PCLP-1

Protein Families: Transmembrane






Product images:




Immunohistochemical staining of paraffinembedded Adenocarcinoma of Human breast tissue using anti-PODXL mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, TA807749) (1:150)




Immunohistochemical staining of paraffinembedded Adenocarcinoma of Human colon tissue using anti-PODXL mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, TA807749) (1:150)






Immunohistochemical staining of paraffinembedded Human liver tissue using anti-PODXL mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, TA807749) (1:150)



Immunohistochemical staining of paraffinembedded Carcinoma of Human liver tissue using anti-PODXL mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, TA807749) (1:150)



Immunohistochemical staining of paraffinembedded Carcinoma of Human thyroid tissue using anti-PODXL mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, TA807749) (1:150)