

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for TA808404

BTBD10 Mouse Monoclonal Antibody [Clone ID: OTI7D3]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI7D3
Applications: IHC, WB

Recommend Dilution: WB 1:2000, IHC 1:150

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human BTBD10(NP_115696) produced in E.coli.

Formulation: PBS (PH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Predicted Protein Size: 53.6 kDa


Gene Name: BTB domain containing 10

Database Link: NP 115696 Entrez Gene 84280 Human


Synonyms: GMRP-1; GMRP1



Product images:




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY BTBD10 ([RC202973], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-BTBD10 (1:2000). Positive lysates [LY410219] (100ug) and [LC410219] (20ug) can be purchased separately from OriGene.




Immunohistochemical staining of paraffinembedded Human liver tissue within the normal limits using anti-BTBD10 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, TA808404) (1:150)





Immunohistochemical staining of paraffinembedded Carcinoma of Human prostate tissue using anti-BTBD10 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, TA808404) (1:150)



Immunohistochemical staining of paraffinembedded Human bladder tissue within the normal limits using anti-BTBD10 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, TA808404) (1:150)