

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## **Product datasheet for TA809230**

## RIAM (APBB1IP) Mouse Monoclonal Antibody [Clone ID: OTI6E3]

**Product data:** 

**Product Type:** Primary Antibodies

Clone Name: OTI6E3
Applications: IHC, WB

Recommend Dilution: WB 1:500~2000, IHC 1:250

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human APBB1IP (NP\_061916) produced in

HEK293T cell.

**Formulation:** PBS (PH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Predicted Protein Size: 73 kDa


**Gene Name:** amyloid beta precursor protein binding family B member 1 interacting protein

Database Link: NP 061916 Entrez Gene 54518 Human


Synonyms: INAG1; PREL1; RARP1; RIAM



## **Product images:**




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY APBB1IP ([RC208712], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-APBB1IP (1:2000). Positive lysates [LY412750] (100ug) and [LC412750] (20ug) can be purchased separately from OriGene.




Western blot analysis of extracts (35ug) from 3 different cell lines by using anti-APBB1IP monoclonal antibody (1:500).





Immunohistochemical staining of paraffinembedded Human Kidney tissue within the normal limits using anti-APBB1IP mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, TA809230) (1:250)



Immunohistochemical staining of paraffinembedded Human lymph node tissue within the normal limits using anti-APBB1IP mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, TA809230) (1:500) (1:250)