

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for TA810146

TSEN34 Mouse Monoclonal Antibody [Clone ID: OTI6B3]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI6B3
Applications: IHC, WB

Recommend Dilution: WB 1:2000, IHC 1:150

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human TSEN34(NP_076980) produced in E.coli.

Formulation: PBS (PH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Predicted Protein Size: 33.5 kDa

Gene Name: tRNA splicing endonuclease subunit 34

Database Link: NP 076980 Entrez Gene 79042 Human

Background: This gene encodes a catalytic subunit of the tRNA splicing endonuclease, which catalyzes the

removal of introns from precursor tRNAs. The endonuclease complex is also associated with a pre-mRNA 3-prime end processing factor. A mutation in this gene results in the neurological disorder pontocerebellar hypoplasia type 2. Multiple alternatively spliced variants, encoding


the same protein, have been identified. [provided by RefSeq, Oct 2009]

Synonyms: LENG5; PCH2C; SEN34; SEN34L






Product images:




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY TSEN34 ([RC200954], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-TSEN34 (1:2000). Positive lysates [LY402979] (100ug) and [LC402979] (20ug) can be purchased separately from OriGene.




Immunohistochemical staining of paraffinembedded Adenocarcinoma of Human colon tissue using anti-TSEN34 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, TA810146) (1:150)






Immunohistochemical staining of paraffinembedded Carcinoma of Human kidney tissue using anti-TSEN34 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, TA810146) (1:150)



Immunohistochemical staining of paraffinembedded Carcinoma of Human lung tissue using anti-TSEN34 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, TA810146) (1:150)



Immunohistochemical staining of paraffinembedded Human endometrium tissue within the normal limits using anti-TSEN34 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, TA810146) (1:150)